The Differential Equation Method: evaluation of complicated Feynman diagrams

A.V.Kotikov

Particle Physics Laboratory, Joint Institute for Nuclear Research
141980 Dubna, Russia.

Abstract

We discuss a progress in calculation of Feynman integrals which has been done with the help of the Differential Equation Method and demonstrate the results for a class of two-point two-loop diagrams.

The idea of the Differential Equation Method (see [1]-[3]) (see reviews in [4])is to apply the integration by parts procedure [5] to an internal n-point subgraph of a complicated Feynman diagram and then to represent new complicated diagrams, obtained this way, as derivatives with respect to corresponding masses of the initial diagram.

The integration by parts procedure [5], [1]-[3] for a general n-point (sub)graph with masses of its lines $m_1, m_2, ..., m_n$, line momenta $p_1, p_2 = p_1 - p_{12}, p_n = p_1 - p_{1n}$ and indices $j_1, j_2, .., j_n$, respectively, has the following form:

$$\begin{aligned} 0 &= \int d^D p_1 \frac{\partial}{\partial p_1^\mu} \left\{ p_1^\mu \left(\prod_{i=1}^{n} c_i^{j_i} \right)^{-1} \right\} \\ &= \int d^D p_1 \left(\prod_{i=1}^{n} c_i^{j_i} \right)^{-1} \left[D - 2j_1 \left(1 - \frac{m_1^2}{c_1} \right) - \sum_{i=2}^{n} j_i \left(1 - \frac{m_1^2 + m_i^2 + p_{1i}^2 - c_1}{c_i} \right) \right], \end{aligned} \tag{1}$$

where $c_k = p_k^2 + m_k^2$ are the propagators of n-point (sub)graph.

Because the diagram with the index $(j_i + 1)$ of the propagator c_i may be represented as the derivative (over the mass m_i), Eq.(1) leads to the differential equations (in principle, to partial differential equations) for the initial diagram (having the index j_i, respectively). This approach which is based on the Eq.(1) and allows to construct the (differential) relations between diagrams has been named as Differential Equations Method (DEM). For the most interested cases (where the number of masses is limited) these partial differential equations may be represented through original differential equation[1], which is usually simpler to analyze.

Thus, we have got the differential equations for the initial diagram. The inhomogeneous term contains only more simple diagrams. These simpler diagrams have more trivial topological structure and/or less number of loops [1] and/or ends [2, 3].

Applying the procedure several times, we will able to represent complicated Feynman integrals (FI) and their derivatives (with respect to internal masses) through a set of quite simple well-known diagrams. Then, the results for the complicated FI can be obtained by several integrations of the known results for corresponding simple diagrams [2].

Sometimes it is useful (see [8]) to use external momenta (or some their functions), but not masses as parameters of integration.

Recent progress in calculation of Feynman integrals with help of the DEM.

1. The articles [9] and [10]:

[1] The example of the direct application of the partial differential equation may be found in [6].

[2] In calculations of real processes (essentially in the framework of Standard Model) it is useful to use the relation (1) (at least, at first steps of calculations) to decrease the number of contributed diagrams (see [1]-[3] and [7] and references therein).



Figure 1: Two-loop selfenergy diagrams. Solid lines denote propagators with the mass m; dashed lines denote massless propagators.

a) The set of two-point two-loop FI with one- and two-mass thresholds has been evaluated by DEM (see Fig.1). The results are given on pages 2 and 3 and some of them have been known before (see [9]). The check of the results has been done by Veretin programs (see discussions in [9, 11] and references therein).

b) The set of three-point two-loop FI with one- and two-mass thresholds has been evaluated (the results of some of them have been known before (see [9])) by a combination of DEM and Veretin programs for calculation of first terms of FI small-moment expansion (see discussions in [9, 11] and references therein).

2. The article [12]:

The full set of two-point two-loop onshell master diagrams has been evaluated by DEM. The check of the results has been done by Kalmykov programs (see page 5 and discussions in [12, 13] and references therein).

3. The articles [14]:

The set of three-point and four-point two-loop massless FI has been evaluated.

Here we demonstrate the results of FI displayed on Fig.1.

We introduce the notation for polylogarithmic functions [15]:

$$\mathrm{Li}_a(z) = S_{a-1,1}(z), \quad S_{a+1,b}(z) = \frac{(-1)^{a+b}}{a!\,b!}\int_0^1 \frac{\log^a(t)\log^b(1-zt)}{t}\,dt.$$

We introduce also the following two variables

$$z = \frac{q^2}{m^2}, \quad y = \frac{1-\sqrt{z/(z-4)}}{1+\sqrt{z/(z-4)}}.$$

Then[3]

$$q^2 \cdot I_1 = -\frac{1}{2}\log^2(-z)\log(1-z) - 2\log(-z)\mathrm{Li}_2(z) + 3\mathrm{Li}_3(z) - 6S_{1,2}(z) - \log(1-z)\Big(\zeta_2 + 2\mathrm{Li}_2(z)\Big),$$

$$\begin{aligned} q^2 \cdot I_5 = {} & 2\zeta_2\log(1+z) + 2\log(-z)\mathrm{Li}_2(-z) + \log^2(-z)\log(1+z) + 4\log(1+z)\mathrm{Li}_2(z) \\ & - 2\mathrm{Li}_3(-z) - 2\mathrm{Li}_3(z) + 2S_{1,2}(z^2) - 4S_{1,2}(z) - 4S_{1,2}(-z), \end{aligned}$$

$$q^2 \cdot I_{12} = \mathrm{Li}_3(z) - 6\zeta_3 - \zeta_2\log y - \frac{1}{6}\log^3 y - 4\log y\,\mathrm{Li}_2(y) + 4\mathrm{Li}_3(y) - 3\mathrm{Li}_3(-y) + \frac{1}{3}\mathrm{Li}_3(-y^3),$$

$$q^2 \cdot I_{13} = -6S_{1,2}(z) - 2\log(1-z)\Big(\zeta_2 + \mathrm{Li}_2(z)\Big),$$

$$\begin{aligned} q^2 \cdot I_{14} = {} & \log(2-z)\Big(\log^2(1-z) - 2\log(-z)\log(1-z) - 2\mathrm{Li}_2(z)\Big) - \frac{2}{3}\log^3(1-z) - 2\zeta_2\log(1-z) \\ & + \log(-z)\log^2(1-z) - S_{1,2}\big(1/(1-z)^2\big) + 2S_{1,2}\big(1/(1-z)\big) + 2S_{1,2}\big(-1/(1-z)\big) + \frac{1}{3}\log^3 y \\ & + \log^2 y\,\Big(2\log(1+y^2) - 3\log(1-y+y^2)\Big) - 6\zeta_3 - \mathrm{Li}_3(-y^2) + \frac{2}{3}\mathrm{Li}_3(-y^3) - 6\mathrm{Li}_3(-y) \\ & + 2\log y\,\Big(\mathrm{Li}_2(-y^2) - \mathrm{Li}_2(-y^3) + 3\mathrm{Li}_2(-y)\Big), \end{aligned}$$

$$\begin{aligned} q^2 \cdot I_{15} = {} & 2\mathrm{Li}_3(z) - \log(-z)\,\mathrm{Li}_2(z) + \zeta_2\log(1-z) - \frac{1}{2}\log^2 y\,\Big(8\log(1-y) - 3\log(1-y+y^2)\Big) - 6\zeta_3 \\ & + \frac{1}{6}\log^3 y - \frac{1}{3}\mathrm{Li}_3(-y^3) + 3\mathrm{Li}_3(-y) + 8\mathrm{Li}_3(y) + \log y\,\Big(\mathrm{Li}_2(-y^3) - 3\mathrm{Li}_2(-y) - 8\mathrm{Li}_2(y)\Big), \end{aligned}$$

$$\begin{aligned} q^2 \cdot I_{123} = {} & -\zeta_2\Big(\log(1-z) + \log y\Big) - 6\zeta_3 - \frac{3}{2}\log(1-y+y^2)\,\log^2 y + \mathrm{Li}_3(-y^3) - 9\mathrm{Li}_3(-y) \\ & - 2\log y\,\Big(\mathrm{Li}_2(-y^3) - 3\mathrm{Li}_2(-y)\Big), \end{aligned}$$

$$q^2 \cdot I_{125} = -2\log^2 y\,\log(1-y) - 6\zeta_3 + 6\mathrm{Li}_3(y) - 6\log y\,\mathrm{Li}_2(y),$$

$$\begin{aligned} q^2 \cdot I_{1234} = {} & -6\zeta_3 - 12\mathrm{Li}_3(y) - 24\mathrm{Li}_3(-y) + 8\log y\,\Big(\mathrm{Li}_2(y) + 2\mathrm{Li}_2(-y)\Big) + 2\log^2 y\,\Big(\log(1-y) \\ & + 2\log(1+y)\Big). \end{aligned} \tag{2}$$

Here we demonstrate the results of FI displayed on Fig.2.

We consider here the following master-integrals in Euclidean space-time with dimension $D = 4 - 2\varepsilon$:

[3] We would like to note that the coefficients of expansions of the results (2) in z are very similar (see [16]) to the results for the moments of structure functions of deep inelastic scattering.

$$
\begin{aligned}
\mathbf{ONS}\{\mathcal{IJ}\}(i,j,m) &\equiv K^{-1}\int d^Dk P^{(i)}(k,\mathcal{I}m)\ P^{(j)}(k-p,\mathcal{J}m)\Big|_{p^2=-m^2}\,,\\
\mathbf{J}\{\mathcal{IJK}\}(i,j,k,m) &\equiv K^{-2}\int d^Dk_1 d^Dk_2 P^{(i)}(k_1,\mathcal{I}m)P^{(j)}(k_1-k_2,\mathcal{J}m)\ P^{(k)}(k_2-p,\mathcal{K}m)\Big|_{p^2=-m^2}\,,\\
\mathbf{V}\{\mathcal{IJKL}\}(i,j,k,l,m) &\equiv K^{-2}\int d^Dk_1 d^Dk_2 P^{(i)}(k_2-p,\mathcal{I}m)\\
&\quad\times P^{(j)}(k_1-k_2,\mathcal{J}m)P^{(k)}(k_1,\mathcal{K}m)\ P^{(l)}(k_2,\mathcal{L}m)\Big|_{p^2=-m^2}\,,\\
\mathbf{F}\{\mathcal{ABIJK}\}(a,b,i,j,k,m) &\equiv m^2K^{-2}\int d^Dk_1 d^Dk_2 P^{(a)}(k_1,\mathcal{A}m)P^{(b)}(k_2,\mathcal{B}m)\\
&\quad\times P^{(i)}(k_1-p,\mathcal{I}m)P^{(j)}(k_2-p,\mathcal{J}m)\ P^{(k)}(k_1-k_2,\mathcal{K}m)\Big|_{p^2=-m^2}\,,
\end{aligned}
$$

where

$$
K=\frac{\Gamma(1+\varepsilon)}{(4\pi)^{\frac{D}{2}}\,(m^2)^{\varepsilon}},\quad P^{(l)}(k,m)\equiv\frac{1}{(k^2+m^2)^l},
$$

the normalization factor $1/(2\pi)^D$ for each loop is assumed, and $\mathcal{A},\mathcal{B},\mathcal{I},\mathcal{J},\mathcal{K}=0,1$.

The finite part of most of the F-type master-integrals can be obtained from results of Ref.[9] in the limit $z\to 1$. **F10101** and **F11111** have been calculated in Refs.[17, 18], respectively. Instead of the usually taken **F01101** integral [17, 19] we consider **J111** as master integral. We recall the results of all master integrals for completeness. The last master integral **F00111** has been found in [12].

The finite part of the integrals of V- and J-type can be found in Refs.[20]. The calculation of the ε e (ε^2) parts of master integrals of this type have been performed by DEM.

The results for F-type master-integrals are follows:

$$
\mathbf{F}\{\mathcal{ABIJK}\}(1,1,1,1,1,m)=a_1\zeta(3)+a_2\frac{\pi}{\sqrt{3}}S_2+a_3 i\pi\zeta(2)+\mathcal{O}(\varepsilon),\tag{3}
$$

and the coefficients $\{a_i\}$ are given in Table I:

TABLE I

	F11111	**F00111**	**F10101**	**F10110**	**F01100**	**F00101**	**F10100**	**F00001**
a_1	-1	0	-4	-1	0	-3	-2	-3
a_2	$\frac{9}{2}$	9	$\frac{27}{2}$	9	$\frac{27}{2}$	$\frac{27}{2}$	9	0
a_3	0	0	$\frac{1}{3}$	0	1	1	$\frac{2}{3}$	1

where [21, 17, 15]

$$
S_2=\frac{4}{9\sqrt{3}}\mathrm{Cl}_2\left(\frac{\pi}{3}\right)=0.260434137632\cdots.
$$

Here we used the $m^2-i\varepsilon$ prescription. The results for the remaining master integrals are the following ones:

$$
\begin{aligned}
\mathbf{V}\{\mathcal{IJKL}\}(1,1,1,1,m) &= \frac{1}{2\varepsilon^2}+\frac{1}{\varepsilon}\left(\frac{5}{2}-\frac{\pi}{\sqrt{3}}\right)+\frac{19}{2}+\frac{b_1}{2}\zeta(2)-4\frac{\pi}{\sqrt{3}}-\frac{63}{4}S_2+\frac{\pi}{\sqrt{3}}\ln 3\\
&+\varepsilon\left\{\frac{65}{2}+b_2\zeta(2)-b_3\zeta(3)-12\frac{\pi}{\sqrt{3}}-63S_2+b_4\zeta(2)\ln 3+\frac{9}{4}b_4S_2\frac{\pi}{\sqrt{3}}\right.\\
&\left.+\frac{63}{4}S_2\ln 3+4\frac{\pi}{\sqrt{3}}\ln 3-\frac{1}{2}\frac{\pi}{\sqrt{3}}\ln^2 3-\frac{b_5}{2}\frac{\pi}{\sqrt{3}}\zeta(2)-\frac{21}{2}\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\right\}+\mathcal{O}(\varepsilon^2),
\end{aligned}
$$

where the coefficients $\{b_i\}$ are listed in Table II [4]:

TABLE II					
	b_1	b_2	b_3	b_4	b_5
V1111	-1	-6	$\frac{9}{2}$	4	9
V1001	3	8	$-\frac{3}{2}$	0	21

$$\begin{aligned}
\mathbf{J111}(1,1,1,m) &= -m^2\left(\frac{3}{2\varepsilon^2}+\frac{17}{4\varepsilon}+\frac{59}{8}+\varepsilon\left\{\frac{65}{16}+8\zeta(2)\right\}\right.\\
&- \left.-\varepsilon^2\left\{\frac{1117}{32}-52\zeta(2)+48\zeta(2)\ln 2-28\zeta(3)\right\}+\mathcal{O}(\varepsilon^3)\right), \qquad (4)\\
\mathbf{J011}(1,1,2,m) &= \frac{1-4\varepsilon}{2(1-2\varepsilon)(1-3\varepsilon)}\left(\frac{1}{\varepsilon^2}+2\frac{\pi}{\sqrt{3}}-\frac{2}{3}\zeta(2)\right.\\
&+ \left.\varepsilon\left\{8\frac{\pi}{\sqrt{3}}-\frac{2}{3}\zeta(2)-6\frac{\pi}{\sqrt{3}}\ln 3+\frac{2}{3}\zeta(3)+27S_2\right\}+\mathcal{O}(\varepsilon^2)\right), \qquad (5)\\
\mathbf{J011}(1,1,1,m) &= -\frac{m^2}{2}\frac{4-15\varepsilon}{(1-2\varepsilon)(1-3\varepsilon)(2-3\varepsilon)}\left(\frac{1}{\varepsilon^2}+\frac{3}{2}\frac{\pi}{\sqrt{3}}+\varepsilon\left\{\frac{45}{8}\frac{\pi}{\sqrt{3}}-\frac{9}{2}\frac{\pi}{\sqrt{3}}\ln 3+\frac{81}{4}S_2\right\}\right.\\
&+ \varepsilon^2\left\{12-\zeta(2)-\frac{1863}{16}S_2-\frac{867}{32}\frac{\pi}{\sqrt{3}}+\frac{207}{8}\frac{\pi}{\sqrt{3}}\ln 3+\frac{243}{4}S_2\ln 3\right.\\
&- \left.\left.\frac{27}{4}\frac{\pi}{\sqrt{3}}\ln^2 3-21\frac{\pi}{\sqrt{3}}\zeta(2)-\frac{81}{2}\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\right\}+\mathcal{O}(\varepsilon^3)\right), \qquad (6)\\
\mathbf{ONS11}(1,1,m) &= \frac{1}{1-2\varepsilon}\left[\frac{1}{\varepsilon}-\frac{\pi}{\sqrt{3}}+\varepsilon\left\{\frac{\pi}{\sqrt{3}}\ln 3-9S_2\right\}\right.\\
&+ \left.\varepsilon^2\left\{9S_2\ln 3-\frac{1}{2}\frac{\pi}{\sqrt{3}}\ln^2 3-6\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}-3\frac{\pi}{\sqrt{3}}\zeta(2)\right\}+\mathcal{O}(\varepsilon^3)\right], \qquad (7)
\end{aligned}$$

where [15]

$$\mathrm{Ls}_3(x)=-\int_0^x \ln^2\left|2\sin\frac{\theta}{2}\right|d\theta \quad \text{and} \quad \mathrm{Ls}_3\left(\frac{2\pi}{3}\right)=-2.14476721256949\cdots$$

The above results were checked numerically. Padé approximants were calculated from the small momentum Taylor expansion of the diagrams [23]. The Taylor coefficients were obtained by means of the package [24] with the master integrals taken from [25]. Further we made use of the idea of Broadhurst [26] to apply the FORTRAN program **PSLQ** [27] to express the obtained numerical values in terms of a 'basis' of irrational numbers, which were predicted by DEM.

Let us point out that the numbers we obtained are related to polylogarithms at the sixth root of unity[5] and hence are in the same class of transcendentals as those obtained by Broadhurst [26] in his investigation of three-loop diagrams which correspond to a closure of the two-loop topologies considered here.

Acknowledgments. The author would like to express his thanks to the Organizing Committees of the Research Workshop "Calculations for modern and future colliders" and the XVth International Workshop

[4] The results for the master integral **V1001** had a little error (see [22])

[5] For the results obtained in $1/N$ expansion, however, the arguments of polylogarithms have other values (see [28]).

"High Energy Physics and Quantum Field Theory" and especially to E.E. Boss, V.A. Ilyin, D.I. Kazakov and D.V. Shirkov for the kind invitation, the financial support at such remarkable Conferences, and for fruitful discussions. Author was supported in part by Alexander von Humboldt fellowship.

References

[1] A.V. Kotikov, Phys.Lett. B254 (1991) 185, Mod.Phys.Lett. A6 (1991) 677.

[2] A.V. Kotikov, Phys.Lett. B259 (1991) 314; Mod.Phys.Lett. A6 (1991) 3133.

[3] A.V. Kotikov, Phys.Lett. B267 (1991) 123; B295 (1992) 409(E); Int.J.Mod.Phys. A6 (1992) 1977.

[4] A.V. Kotikov, JHEP 9809 (1998) 001; in: AIHENP 93, *Proceedings of the 3th International Workshop on Software Engineering, Artificial Intelligence and Expert Systems*, ed.by D. Perret-Gallix, in pp. 539-544; in: ACAT 2000, *Proceedings of the 7th International Workshop on Software Engineering, Artificial Intelligence and Expert Systems*, ed.by D. Perret-Gallix, in American Institute of Physics Press., (hep-ph/0011316).

[5] F.V. Tkachov, Phys.Lett. B100 (1981) 65; K.G. Chetyrkin and F.V. Tkachov, Nucl.Phys. B192 (1981) 159; A.N. Vassiliev, Yu.M. Pis'mak and Yu.R. Khonkonen, Theor.Math.Phys. B47 (1981) 291.

[6] C. Ford, I. Jack and D.R.T. Jones, Nucl.Phys. B387 (1992) 373.

[7] J. Fleischer, M. Tentyukov, and O.V. Tarasov, Nucl.Phys.Proc.Suppl. 89 (2000) 112.

[8] E. Remiddi, Nuovo Cim. A110 (1997) 1435.

[9] J. Fleischer, A.V. Kotikov, and O.L. Veretin, Nucl.Phys. B547 (1999) 343.

[10] J. Fleischer, A.V. Kotikov, and O.L. Veretin, Phys.Lett. B417 (1998) 163.

[11] J. Fleischer, A.V. Kotikov, and O.L. Veretin, Acta Phys.Polon. B29 (1998) 2611.

[12] J. Fleischer, M.Yu. Kalmykov, and A.V. Kotikov, Phys.Lett. B462 (1999) 169; B467 (1999) 310(E).

[13] J. Fleischer, M.Yu. Kalmykov, and A.V. Kotikov, in AIHENP 99, *Proceedings of the 6th International Workshop on Software Engineering, Artificial Intelligence and Expert Systems*, ed.by G. Athanasiu and D. Perret-Gallix, in Parisianou S.A., Athens, 2000, pp. 231-237, (hep-ph/0011316); J. Fleischer and M.Yu. Kalmykov, Comput.Phys.Commun. 128 (2000) 531.

[14] T. Gehrmann and E. Remiddi, Nucl.Phys. B580 (2000) 485; Preprint TTP00-20, 2000 (hep-ph/0008287); Preprint CERN-TH/2001-005, 2001 (hep-ph/0101124); Preprint TTP01-04, 2001 (hep-ph/0101147); Nucl.Phys.Proc.Suppl. 89 (2000) 251; C. Anastasiou, T. Gehrmann, C. Oleari, E. Remiddi and J.B. Tausk, Nucl.Phys. B580 (2000) 577.

[15] L. Lewin, Polylogarithms and Associated Functions, North Holland, New-York, 1981.

[16] D.I. Kazakov and A.V. Kotikov, Theor.Math.Phys. 73 (1987) 1264; Nucl.Phys. B307 (1988) 721; B345 (1990) 299(E); Phys.Lett. B291 (1992) 171; A.V. Kotikov, Theor.Math.Phys. 78 (1989) 134.

[17] D.J. Broadhurst, Z.Phys. C47 (1990) 115.

[18] V. Borodulin and G. Jikia, Phys.Lett. B391 (1997) 434.

[19] N. Gray, D.J. Broadhurst, W. Grafe and K. Schilcher, Z.Phys. C48 (1990) 673; D.J. Broadhurst, N. Gray and K. Schilcher, Z.Phys. C52 (1991) 111; D.J. Broadhurst, Z.Phys. C54 (1992) 599.

[20] A. Djouadi, Nuovo Cim. A100 (1988) 357; P.N. Maher, L. Durand and K.Riesselmann, Phys.Rev. D48 (1993) 1061; D52 (1995) 553(E); R. Scharf and J.B. Tausk, Nucl.Phys. B412 (1994) 523; F.A. Berends, M. Buza, M. Böhm and R. Scharf, Z.Phys. C63 (1994) 227; F.A. Berends and J.B. Tausk, Nucl.Phys. B421 (1994) 456; S. Bauberger, F.A. Berends, M. Böhm and M. Buza, Nucl.Phys. B434 (1995) 383.

[21] J. van der Bij and M. Veltman, Nucl.Phys. B231 (1984) 205; F. Hoogeveen, Nucl.Phys. B259 (1985) 19; J. van der Bij and F. Hoogeveen, Nucl.Phys. B283 (1987) 477.

[22] A.I. Davydychev and M.Yu. Kalmykov, Preprint MZ-TH/00-52, 2000, (hep-th/0012189).

[23] J. Fleischer and O.V. Tarasov, Z.Phys. C64 (1994) 413; O.V. Tarasov, Nucl.Phys. B480 (1996) 397.

[24] L.V. Avdeev, J. Fleischer, M.Yu. Kalmykov and M.N. Tentyukov, Nucl.Inst.Meth. A389 (1997) 343; Comp.Phys.Commun. 107 (1997) 155.

[25] A.I. Davydychev and J.B. Tausk, Nucl.Phys. B397 (1993) 123.

[26] D.J. Broadhurst, Eur.Phys.J. C8 (1999) 311.

[27] H.R.P. Ferguson, D.H. Bailey and S. Arno, NASA-Ames Technical Report, NAS-96-005; D.H. Bailey and D.J. Broadhurst, math.NA/9905048.

[28] D.J. Broadhurst, J.A. Gracey, and D. Kreimer, Z.Phys. C75 (1997) 559; D.J. Broadhurst and A.V. Kotikov, Phys.Lett. B441 (1998) 345; A.V. Kotikov and L.N. Lipatov, Nucl.Phys. B582 (2000) 19.



Figure 2: The full set of two-loop self-energy diagrams with one mass. Bold and thin lines correspond to massive and massless propagators respectively.