

Chains with the fractal dispersion law

Vasily E Tarasov

Skobeltsyn Institute of Nuclear Physics, Moscow State University, Moscow 119991, Russia

E-mail: tarasov@theory.sinp.msu.ru

Received 6 October 2007, in final form 14 November 2007

Published 4 January 2008

Online at stacks.iop.org/JPhysA/41/035101

Abstract

Chains with long-range interactions are considered. The interactions are defined such that each n th particle interacts only with chain particles with the numbers $n \pm a(m)$, where $m = 1, 2, 3, \dots$ and $a(m)$ is an integer-valued function. Exponential type functions $a(m) = b^m$, where $b = 2, 3, \dots$, are discussed. The correspondent pseudodifferential equations of chain oscillations are obtained. Dispersion laws of the suggested chains are described by the Weierstrass and Weierstrass–Mandelbrot functions.

PACS number: 05.45.Df; 45.05.+x; 45.50.-j

1. Introduction

Long-range interaction (LRI) has been the subject of investigations for a long time. An infinite Ising chain with LRI was considered by Dyson [1]. The d -dimensional Heisenberg model with long-range interaction is described in [2, 3], and its quantum generalization can be found in [4, 5]. Solitons in a chain with the long-range Lennard–Jones-type interaction were considered in [7]. Kinks in the Frenkel–Kontorova chain model with long-range interparticle interactions were studied in [8]. The properties of time periodic spatially localized solutions (breathers) on discrete chains in the presence of algebraically decaying LRI were described in [9]. Energy and decay properties of discrete breathers in chains with LRI have also been studied in the framework of the Klein–Gordon [6] and discrete nonlinear Schrödinger equations [10]. A main property of the chain dynamics with power-like long-range interactions [9, 11] is that the solutions of chain equations have power-like tails. The power-law LRI is also considered in [12–15]. In [24], we formulate the consistent definition of continuous limit for the chains with long-range interactions (LRI). In the continuous limit, the chain equations with power-like LRI give the medium equations with fractional derivatives.

Usually we assume for LRI that each chain particle acts on all chain particles. There are systems where this assumption cannot be used. In general, the chain cannot be considered as a straight line. For example, the linear polymers can be represented as some compact objects. It is well known that ‘tertiary structure’ of proteins refers to the overall folding of the entire

polypeptide chain into a specific 3D shape [16–18]. The tertiary structure of enzymes is often compact, globular shaped [16, 17]. In this case, we can consider that the chain particle is interacted with particles of a ball with radius R . Then only some subset A_n of chain particles act on n th particle. We suppose that n th particle is interacted only with k th particles with $k = n \pm a(m)$, where $a(m) \in \mathbb{N}$ and $m = 1, 2, 3, \dots$. We can consider fractal compactified linear polymers (chains), such that these ‘compact objects’ satisfy the power law $N(R) \sim R^D$, where $2 < D < 3$ and $N(R)$ is the number of chain particles inside the sphere with radius R . As an example of such case, $a(m)$ will be described by exponential type functions $a(m) = b^m$, where $b > 1$ and $b \in \mathbb{N}$. In this case, the LRI will be called fractal interaction.

The goal of this paper is to study a connection between the dynamics of chain with fractal long-range interactions (FLRI) and the continuous medium equations with fractal dispersion law. Here, we consider the chain of coupled linear oscillators with FLRI. We make the transformation to the continuous field and derive the continuous equation which describes the dynamics of the oscillatory medium. We show how the oscillations of chains with FLRI are described by the fractal dispersion law. This law is represented by the Weierstrass functions whose graphs have non-integer box-counting dimension, i.e. these graphs are fractals. Fractals are good models of phenomena and objects in various areas of science [20]. Note that fractals in quantum theory have recently been considered in [21]. In this paper, we prove that the chains with long-range interaction can demonstrate fractal properties described by fractal functions.

2. Chain equation

One of the oldest fractal functions is the Weierstrass function [22]:

$$W(x) = \sum_{n=0}^{\infty} b^n \cos(a^n \pi x) \tag{1}$$

introduced as an example of everywhere continuous nowhere differentiable function by Karl Weierstrass around 1872. Maximum range of parameters for which the above sum has fractal properties was found by Godfrey Harold Hardy [23] in 1916, who showed that

$$0 < b < 1, \quad ab \geq 1.$$

The box-counting dimension of the graph of the Weierstrass function $W(x)$ is

$$D = 2 + \frac{\ln(b)}{\ln(a)} = 2 - \left| \frac{\ln(b)}{\ln(a)} \right|. \tag{2}$$

Functions whose graphs have non-integer box-counting dimension are called fractal functions.

Consider a one-dimensional system of interacting oscillators that are described by the equations of motion,

$$\frac{d^2}{dt^2} u_n(t) = c^2 \sum_{m=-\infty}^{+\infty} \frac{b(m)}{h^2} [u_{n+a(m)}(t) - 2u_n(t) + u_{n-a(m)}(t)], \tag{3}$$

where $u_n(t)$ are displacements from the equilibrium and h is the distance between the oscillators. Here $a(m)$ and $b(m)$ are some functions of integer number m . The case $b(m) = J(|n - m|)$ has been considered in [24]. The right-hand side of the equation describes an interaction of the oscillators in the system.

We illustrate this chain equation with well-known example [26]. In the case of nearest-neighbor interaction, we have

$$a(m) = 1, \quad b(m) = \delta m_0.$$

Then equation (3) gives

$$\frac{d^2}{dt^2} u_n(t) = \frac{c^2}{h^2} [u_{n+1}(t) - 2u_n(t) + u_{n-1}(t)]. \quad (4)$$

We can define a smooth function $u(x, t)$ such that

$$u(nh, t) = u_n(t). \quad (5)$$

Then equation (4) has the form

$$\partial_t^2 u(x, t) = \frac{c^2}{h^2} [u(x + h, t) - 2u(x, t) + u(x - h, t)]. \quad (6)$$

This is the differential-difference equation. Using the relation

$$\exp(-ih\partial_x)u(x, t) = u(x + h, t),$$

we can rewrite equation (6) in the form

$$\partial_t^2 u(x, t) + \frac{4c^2}{h^2} \sin^2\left(-\frac{ih}{2}\partial_x\right)u(x, t) = 0. \quad (7)$$

This is the pseudo-differential equation. The properties of this equation have been considered in [26]. For $h \rightarrow 0$, we obtain the wave equation

$$\partial_t^2 u(x, t) - c^2 \partial_x^2 u(x, t) = 0.$$

3. Fractal long-range interaction

If $a(m)$ in equation (1) is not a constant function, then we have the long-range interaction of the chain particles. Note that the function $a(m)$ should be integer valued. For example, $a(m) = m$ and $a(m) = 2^m$. The set $A_n = \{n \pm a(m) : m \in \mathbb{N}\}$ describes the numbers of particles that act on the n th particle.

(a) If $a(m) = m$, where $m \in \mathbb{N}$, then A_n is a set of all integer numbers \mathbb{Z} for all n , i.e., $A_n = \mathbb{Z}$. In this case, the n th particle interacts with all chain particles.

(b) If $a(m) = 2^m$, where $m \in \mathbb{N}$, then A_n is a subset of \mathbb{Z} , i.e., $A_n \subset \mathbb{Z}$. In this case, the n th particle interacts only with chain particles with numbers $n \pm 2, n \pm 4, n \pm 8, n \pm 16 \dots$

The power law $a(m) = b^m$, where $b \in \mathbb{N}$, and $b > 1$, can be realized for compact structure of linear polymer molecules. For example, a linear polymer molecule is not a straight line. Usually this molecule can be considered as a compact object. It is well known that ‘tertiary structure’ of proteins refers to the overall folding of the entire polypeptide chain into a specific 3D shape [16–18]. The tertiary structure of enzymes is often a compact, globular shape [16, 17]. In this case, we can consider that the chain particle is interacted with particles inside a sphere with radius R . Then only some subset of chain particles act on n th particle, We assume that n th particle is interacted only with k th particles with $k = n \pm a(m)$, where $a(m) \in \mathbb{N}$ and $m = 1, 2, 3, \dots$. The polymer can be a mass fractal object [19]. For fractal compactified linear polymer chains, we have the power law $N(R) \sim R^D$, where $2 < D < 3$ and $N(R)$ is the number of chain particles in the ball with radius R . Then we suppose that $a(m)$ is exponential type function such that $a(m) = b^m$, where $b > 1$ and $b \in \mathbb{N}$. This function defines the fractal long-range interaction.

Let us consider equation (3) with a fractal long-range interaction. Using the smooth function (5), we obtain the differential-difference equation

$$\partial_t^2 u(x, t) = \frac{c^2}{h^2} \sum_{m=-\infty}^{+\infty} b(m) [u(x + a(m)h, t) - 2u(x, t) + u(x - a(m)h, t)]. \quad (8)$$

Using

$$\exp(-ia(m)h\partial_x)u(x, t) = u(x + a(m)h, t), \tag{9}$$

equation (8) can be presented as the pseudo-differential equation

$$\partial_t^2 u(x, t) + \frac{4c^2}{h^2} \sum_{m=-\infty}^{+\infty} b(m) \sin^2\left(-\frac{iha(m)}{2}\partial_x\right)u(x, t) = 0. \tag{10}$$

For $a(m) = 1$ and $b(m) = \delta m_0$, this equation gives equation (7) that describes oscillations for the case of the nearest-neighbor interaction.

Let us consider the pseudo-differential operator

$$\mathcal{L} = 2 \sum_{m=-\infty}^{+\infty} b(m) \sin^2\left(-\frac{iha(m)}{2}\partial_x\right). \tag{11}$$

The function

$$\Psi(x, k) = A \exp(ikx) \tag{12}$$

is an eigenfunction of this operator:

$$\mathcal{L}\Psi(x, k) = \lambda(k)\Psi(x, k).$$

Here $\lambda(k)$ is the eigenvalue of the operator, such that

$$\lambda(k) = 2 \sum_{m=-\infty}^{+\infty} b(m) \sin^2\left(\frac{ha(m)}{2}k\right).$$

Using $2 \sin^2(\alpha/2) = 1 - \cos(\alpha)$, we obtain

$$\lambda(k) = \sum_{m=-\infty}^{+\infty} b(m)[1 - \cos(ha(m)k)]. \tag{13}$$

If

$$a(m) = a^m, \quad b(m) = a^{(D-2)m}, \quad a \in \mathbb{N}, \quad a > 1, \tag{14}$$

then equation (13) has the form

$$\lambda(k) = C(hk),$$

where $C(z)$ is the cosine Weierstrass–Mandelbrot function [20, 25],

$$C(z) = \sum_{m=-\infty}^{+\infty} a^{(D-2)m}[1 - \cos(a^{(D-2)m}z)].$$

The box-counting dimension of the graph of this function is D . The operator (11) can be called the Weierstrass–Mandelbrot operator. The spectral graph $(k, C(hk))$ of this operator is a fractal set with dimension D .

Using the operator (11), equation (10) takes the form

$$\partial_t^2 u(x, t) + \frac{4c^2}{h^2} \mathcal{L}u(x, t) = 0.$$

It is not hard to prove that the dispersion law for the chain with the long-range interactions is described by equations (3) and (14) has the form

$$\omega^2 + \frac{2c^2}{h^2} C(hk) = 0.$$

Then the graph $(k, \omega(k))$ is a fractal. Note that the group velocity $v_{\text{group}} = \partial\omega(k)/\partial k$ for the plane waves cannot be found, since $C(z)$ is the nowhere differentiable function.

Let us consider a generalization of conditions (14) in the form

$$a(m) = a^m, \quad b(m) = b^m, \quad (15)$$

where $a \in \mathbb{N}$, $b < 1$ and $a > 1$. For $b = a^{D-2}$, we have equation (14). If we use equation (15), we can obtain the pseudo-differential equation

$$\partial_t^2 u(x, t) + \frac{4c^2}{h^2} \sin^2\left(-\frac{ih}{2}\partial_x\right)u(x, t) + \frac{8c^2}{h^2} \sum_{m=1}^{+\infty} b^m \sin^2\left(-\frac{iha^m}{2}\partial_x\right)u(x, t). \quad (16)$$

This equation can be presented in the form

$$\partial_t^2 u(x, t) + \frac{4c^2}{h^2} \sin^2\left(-\frac{ih}{2}\partial_x\right)u(x, t) + M^2 c^2 u(x, t) = 4\frac{c^2}{h^2} \sum_{m=1}^{+\infty} b^m \cos\left(-iha^m\partial_x\right)u(x, t), \quad (17)$$

where $M^2 = 4b/h^2(1 - b)$. Equation (17) describes the oscillations in the case of the fractal LRI. The left-hand side of equation (17) in the limit $h \rightarrow 0$ gives the Klein–Gordon equation

$$c^{-2}\partial_t^2 u(x, t) - \partial_x^2 u(x, t) + M^2 u(x, t) = 0.$$

The right-hand side of equation (17) describes a nonlocal part of the interaction. Note that the pseudo-differential operator

$$\Lambda = \sum_{m=0}^{+\infty} b^m \cos(-iha^m\partial_x), \quad (18)$$

which is used in equation (17), has the eigenfunctions (12) such that the eigenvalues is the Weierstrass function $W(hk/\pi)$, where $W(x)$ is defined by equation (1). The box-counting dimension of the graph of the Weierstrass function $W(x)$ is (2). The operator (18) can be called the Weierstrass operator. The spectral graph $(k, W(hk/\pi))$ of this operator is a fractal set with dimension $2 + \ln(b)/\ln(a)$.

4. Conclusion

In this paper, we prove that the chains with long-range interaction can demonstrate fractal properties. We consider chains with long-range interactions such that each n th particle is interacted only with chain particles with the numbers $n \pm a(m)$, where $m = 1, 2, 3, \dots$. The exponential type functions $a(m) = b^m$, where $b > 1$ is integer, are used to define a fractal long-range interaction. The equations of chain oscillations are characterized by dispersion laws that are represented by Weierstrass and Weierstrass–Mandelbrot (fractal) functions. The suggested chains with long-range interactions can be considered as a simple model for linear polymers that are compact, fractal globular shaped.

References

- [1] Dyson F J 1969 Existence of a phase-transition in a one-dimensional Ising ferromagnet *Commun. Math. Phys.* **12** 91–107
- Dyson F J 1969 Non-existence of spontaneous magnetization in a one-dimensional Ising ferromagnet *Commun. Math. Phys.* **12** 212–5
- Dyson F J 1971 An Ising ferromagnet with discontinuous long-range order *Commun. Math. Phys.* **21** 269–83

- [2] Joyce G S 1969 Absence of ferromagnetism or antiferromagnetism in the isotropic Heisenberg model with long-range interactions *J. Phys. C: Solid State Phys.* **2** 1531–3
- [3] Fröhlich J, Israel R, Lieb E H and Simon B 1978 Phase transitions and reflection positivity I. General theory and long-range lattice model *Comm. Math. Phys.* **62** 1–34
- [4] Nakano H and Takahashi M 1994 Quantum Heisenberg model with long-range ferromagnetic interactions *Phys. Rev. B* **50** 10331–4
Nakano H and Takahashi M 1995 Magnetic properties of quantum Heisenberg ferromagnets with long-range interactions *Phys. Rev. B* **52** 6606–10
- [5] Sousa J R 2005 Phase diagram in the quantum XY model with long-range interactions *Eur. Phys. J. B* **43** 93–6
- [6] Braun O M and Kivshar Y S 1998 Nonlinear dynamics of the Frenkel–Kontorova model *Phys. Rep.* **306** 2–108
- [7] Ishimori Y 1982 Solitons in a one-dimensional Lennard–Jones lattice *Prog. Theor. Phys.* **68** 402–10
- [8] Braun O M, Kivshar Y S and Zelenskaya I 1990 Kinks in the Frenkel–Kontorova model with long-range interparticle interactions *Phys. Rev. B* **41** 7118–38
- [9] Flach S 1998 Breathers on lattices with long-range interaction *Phys. Rev. E* **58** R4116–9
Gorbach A V and Flach S 2005 Compactlike discrete breathers in systems with nonlinear and nonlocal dispersive terms *Phys. Rev. E* **72** 056607
- [10] Gaididei Yu B, Mingaleev S F, Christiansen P L and Rasmussen K O 1997 Effects of nonlocal dispersive interactions on self-trapping excitations *Phys. Rev. E* **55** 6141–50
Mingaleev S F, Gaididei Y B and Mertens F G 1998 Solitons in anharmonic chains with power-law long-range interactions *Phys. Rev. E* **58** 3833–42
Mingaleev S F, Gaididei Y B and Mertens F G 2000 Solitons in anharmonic chains with ultra-long-range interatomic interactions *Phys. Rev. E* **61** R1044–7
Rasmussen K O, Christiansen P L, Johansson M, Gaididei Yu B and Mingaleev S F 1998 Localized excitations in discrete nonlinear Schrödinger systems: effects of nonlocal dispersive interactions and noise *Physica D* **113** 134–51
- [11] Alfimov G L, Eleonsky V M and Lerman L M 1998 Solitary wave solutions of nonlocal sine-gordon equations *Chaos* **8** 257–71
Alfimov G L and Korolev V G 1998 On multikink states described by the nonlocal sine-gordon equation *Phys. Lett. A* **246** 429–35
- [12] Laskin N and Zaslavsky G M 2006 Nonlinear fractional dynamics on a lattice with long-range interactions *Physica A* **368** 38–54
- [13] Tarasov V E and Zaslavsky G M 2006 Fractional dynamics of coupled oscillators with long-range interaction *Chaos* **16** 023110
Tarasov V E and Zaslavsky G M 2006 Fractional dynamics of systems with long-range interaction *Commun. Nonlinear Sci. Numer. Simul.* **11** 885–98
- [14] Korabel N, Zaslavsky G M and Tarasov V E 2007 Coupled oscillators with power-law interaction and their fractional dynamics analogues *Commun. Nonlinear. Sci. Numer. Simul.* **12** 1405–17
- [15] Korabel N and Zaslavsky G M 2007 Transition to chaos in discrete nonlinear Schrödinger equation with long-range interaction *Physica A* **378** 223–37
- [16] Holde K E van 1998 *Principles of Physical Biochemistry* (Englewood Cliffs, NJ: Prentice-Hall)
- [17] The RCSB Protein Data Bank (PDB): <http://www.rcsb.org/pdb/home/home.do>
- [18] Kolinski A and Skolnick J 2004 Reduced models of proteins and their applications *Polymer* **45** 511–24
- [19] Newkome G R *et al* 2006 Nanoassembly of a fractal polymer: a molecular Sierpinski hexagonal gasket *Science* **312** 1782–5
- [20] Mandelbrot B *The Fractal Geometry of Nature* (San Francisco: Freeman)
- [21] Kröger H 2000 Fractal geometry in quantum mechanics, field theory and spin systems *Phys. Rep.* **323** 818–181
Berry M V 1996 Quantum fractals in boxes *J. Phys. A: Math. Gen.* **29** 6617–29
Wojcik D, Białynicki-Birula I and Zyczkowski K 2000 Time evolution of quantum fractals *Phys. Rev. Lett.* **85** 5022–6
- [22] Weierstrass F Über kontinuierliche funktionen eines reellen arguments, die für keinen wert des letzteren einen bestimmten differentialquotienten besitzen In *Mathematische Werke II* (Berlin: Mayer-Müller) 71–4
- [23] Hardy G H 1916 Weierstrass's non-differentiable function *Trans. AMS* **17** 301–25
- [24] Tarasov V E 2006 Continuous limit of discrete systems with long-range interaction *J. Phys. A* **39** 14895–910
- [25] Feder J *Fractals* (New York: Plenum)
- [26] Maslov V P *Operator Method* (Moscow: Nauka) (in Russian)
Maslov V P *Operational Methods* (Moscow: Mir) (English transl.)